



**Cochrane**  
**Library**

Cochrane Database of Systematic Reviews

## Selenium for alleviating the side effects of chemotherapy, radiotherapy and surgery in cancer patients (Review)

Dennert G, Horneber M

Dennert G, Horneber M.

Selenium for alleviating the side effects of chemotherapy, radiotherapy and surgery in cancer patients.

*Cochrane Database of Systematic Reviews* 2006, Issue 3. Art. No.: CD005037.

DOI: 10.1002/14651858.CD005037.pub2.

[www.cochranelibrary.com](http://www.cochranelibrary.com)

## TABLE OF CONTENTS

HEADER . . . . .	1
ABSTRACT . . . . .	1
PLAIN LANGUAGE SUMMARY . . . . .	2
BACKGROUND . . . . .	2
OBJECTIVES . . . . .	3
METHODS . . . . .	3
RESULTS . . . . .	5
DISCUSSION . . . . .	7
AUTHORS' CONCLUSIONS . . . . .	9
ACKNOWLEDGEMENTS . . . . .	9
REFERENCES . . . . .	9
CHARACTERISTICS OF STUDIES . . . . .	14
DATA AND ANALYSES . . . . .	20
ADDITIONAL TABLES . . . . .	20
APPENDICES . . . . .	23
WHAT'S NEW . . . . .	24
HISTORY . . . . .	25
CONTRIBUTIONS OF AUTHORS . . . . .	25
DECLARATIONS OF INTEREST . . . . .	25
SOURCES OF SUPPORT . . . . .	25
NOTES . . . . .	26
INDEX TERMS . . . . .	26

[Intervention Review]

# Selenium for alleviating the side effects of chemotherapy, radiotherapy and surgery in cancer patients

Gabriele Dennert<sup>1</sup>, Markus Horneber<sup>2</sup>

<sup>1</sup>Social Medicine and Public Health with Focus on Gender and Diversity, Department of Applied Social Sciences, University of Applied Sciences Dortmund, Dortmund, Germany. <sup>2</sup>Department of Internal Medicine, Division of Oncology and Hematology, Paracelsus Medical University, Klinikum Nuremberg, Nuremberg, Germany

Contact address: Gabriele Dennert, Social Medicine and Public Health with Focus on Gender and Diversity, Department of Applied Social Sciences, University of Applied Sciences Dortmund, Emil-Figge-Str. 44, Dortmund, D-44227, Germany. [gabriele.dennert@fh-dortmund.de](mailto:gabriele.dennert@fh-dortmund.de).

**Editorial group:** Cochrane Pain, Palliative and Supportive Care Group.

**Publication status and date:** Stable (no update expected for reasons given in 'What's new'), published in Issue 3, 2017.

**Citation:** Dennert G, Horneber M. Selenium for alleviating the side effects of chemotherapy, radiotherapy and surgery in cancer patients. *Cochrane Database of Systematic Reviews* 2006, Issue 3. Art. No.: CD005037. DOI: 10.1002/14651858.CD005037.pub2.

Copyright © 2017 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

## ABSTRACT

### Background

This is an updated version of the original Cochrane review published in Issue 3, 2006. Selenium supplements are frequently used by cancer patients. Selenium is an essential trace element and is involved in antioxidant protection and the redox-regulation in humans. Several adverse effects of radiotherapy and chemotherapy in cancer patients as well as cellular processes that maintain chronic lymphoedema have been linked to oxidative cell processes in the human body. Selenium has been claimed to alleviate side effects of conventional cancer therapy and recently been investigated as a remedy against chemotherapy and radiotherapy-associated side effects and secondary lymphoedema.

### Objectives

This review assessed the effects of supplementary selenium on adverse effects of conventional radiotherapy, chemotherapy, or surgery in oncology patients and on quality of life/performance status during and after oncological treatment.

### Search methods

The Cochrane Pain, Palliative & Supportive Care Trials Register, the Cochrane Database of Systematic Reviews, the Database of Abstracts of Reviews of Effects (DARE), CENTRAL, MEDLINE, EMBASE, SIGLE, Cancerlit, CCMed, the German online register for cancer trials, the ISRCTN, the mRCT, the NCI Clinical Trials register databases were searched and most updated in July 2007.

### Selection criteria

Randomised controlled trials (RCTs) of selenium mono-supplements in cancer patients undergoing tumour specific therapy, such as chemotherapy, radiotherapy or surgery.

### Data collection and analysis

Two review authors independently checked trials for eligibility, extracted data and assessed trial quality.

## Main results

One new included study as well as further participants to a study already included was added to this update involving a further 39 participants. There were a total of three studies included involving 162 participants for this update.

For this review two RCTs were included investigating secondary lymphoedema in 60 women after breast cancer surgery and 20 men and women after head and neck cancer surgery. One ongoing trial with preliminary results of 82 participants was also identified studying radiotherapy induced diarrhoea as a secondary outcome. All studies had considerable drawbacks with regards to quality and reporting.

One study on secondary lymphoedema reported a decreased number of recurrent erysipela infections in the selenium supplementation group compared to placebo. The second study reported a decreased facial swelling in the selenium group in a two-week period following surgical tumour resection. However, results must be interpreted with caution and cannot be generalised to other populations.

The ongoing trial on radiotherapy associated diarrhoea preliminarily reported a lower incidence of diarrhoea in participants receiving selenium supplementation concomitant to pelvic radiation, however, no data were presented. We must await publication of final results to discuss these findings in detail.

No RCTs were found studying the effect of selenium supplementation on other therapy-associated toxicities or quality of life/performance status in cancer patients.

## Authors' conclusions

There is insufficient evidence at present that selenium supplementation alleviates the side effects of tumour specific chemotherapy or radiotherapy treatments or that it improves the after-effects of surgery, or improves quality-of-life in cancer patients or reduces secondary lymphoedema. To date, research findings do not provide a basis for any recommendation in favour or against selenium supplementation in cancer patients. Potential hazards of supplementing a trace mineral should be kept in mind. Since the last version of this review, the one new additional study has not provided information to change the conclusions of the original review.

## PLAIN LANGUAGE SUMMARY

### Selenium supplements for alleviating the side effects of chemotherapy, radiotherapy and surgery in cancer patients

There is no evidence that selenium supplements reduce side effects of chemotherapy, radiotherapy or the effects of surgery in cancer patients. Selenium is a mineral necessary for human health. Selenium acts against cell damage in the body and might help to alleviate the side effects of therapies in cancer patients, such as nausea, diarrhoea or the lymph retention in limbs. Selenium supplements are frequently used by cancer patients. This systematic review looked at studies providing selenium supplements to cancer patients and found no clear evidence that selenium supplements improve side effects of cancer therapy. No adverse effects were reported in the studies, but evidence of overdosing, all be it unintentional and selenium intoxication has occurred in several selenium users. More research is needed to find out which doses of selenium supplements can be reasonably used by cancer patients and whether selenium supplements can affect the side effects of cancer therapy.

## BACKGROUND

This review is an update of a previously published review in The Cochrane Database of Systematic Reviews (Issue 3, 2006) on 'Selenium for alleviating the side effects of chemotherapy, radiotherapy and surgery in cancer patients'.

Selenium is an essential trace element for humans. It is involved in antioxidant protection and redox regulation as a component

of seleno-proteins and as a source of seleno-metabolites (Hatfield 2001). Adverse effects of radiotherapy and chemotherapy in cancer patients have been linked to the formation of free radicals and the related oxidative damage to normal cells (Weijl 1997). Lately, intervention strategies have been investigated to use biological response modifiers, such as selenium compounds, as toxicity antagonists for prevention of chemotherapy and radiotherapy associated

side effects.

Acute mucositis is a severe condition seen frequently in patients undergoing radiotherapy. A recent review found that about 80% of ENT radiotherapy patients developed mucositis, about half of them experienced a severe form (grade three to four) which lead to treatment interruption or alteration in one in nine patients (Trotti 2003). Patients undergoing pelvic radiation for e.g. gynecological malignancies are at risk for developing acute intestinal mucositis, enteritis, and diarrhoea. Atrophy of the salivary glands and xerostomia are long-term sequelae of a radiogenic sialadenitis after ENT radiotherapy, which can only be controlled insufficiently by symptomatic interventions, such as saliva substitutes or stimulants (Groetz 2003).

Both in acute and long-term side effects of radiotherapy, selenium has been discussed as a preventive agent. In vitro experiments found a cytoprotective effect of sodium selenite on human fibroblasts and endothelial cells without reducing radiotherapy activity against cancer cells (Rodemann 1999; Schleicher 1999). Sagowski 2004 showed that administration of parenteral sodium selenite reduced radiogenic damage to parotid glands in rats, which was confirmed by better gland function after irradiation.

A number of chemotherapy toxicities and adverse effects have been linked to the formation of free radicals by cytostatics (Weijl 1997). A protective effect of selenium against doxorubicin induced cell damage was seen in rat cardiomyocytes in vitro (Kotamraju 2000). Animal studies suggested that sodium selenite may decrease cisplatin induced nephrotoxicity (Baldew 1989; Francescato 2001; Yoshida 2000) and myelotoxicity (Ohkawa 1988) in rats and mice. Cardiac sensitivity to ischemia after adriamycin treatment was reduced by sodium selenite supplementation in rats (Boucher 1995).

Secondary lymphoedema is a common complication after surgical and radiological therapy of breast and ENT cancer (Dietz 1998; Erickson 2001). Patients with lymphoedema of the upper extremity and the head and neck region can experience a substantial degree of functional impairment and psychological morbidity. Endolaryngeal oedema and swelling may even lead to airway obstruction and require tracheostomy. The development of oedema is indicated by surgical or radiological damage to the lymphatic system resulting in fluid retention (Zimmermann 2005). Higher interstitial pressure lessens oxygen supply to oedematous tissue and chronic inflammatory processes lead to fibrosis of small lymph vessels. Highly reactive oxygen containing radicals (reactive oxygen species) are claimed to have a crucial role in development and maintenance of lymphoedema. Selenium compounds may improve redox balance in sparsely perfused oedematous tissue and, hence, be effective in control and therapy of secondary lymphoedema (Micke 2003; Zimmermann 2005).

The role of the complementary use of selenium in cancer patients is unclear. Some publications report a positive effect of selenium supplementation on radio/chemotherapy associated side effects

and lymphoedema following surgical cancer treatment. However, trial evidence is inconsistent, and a systematic review on this topic has not been conducted.

Surveys in Canada, UK, Austria and Germany found that four to 12% of breast cancer and prostate cancer patients used selenium supplements during and after cancer therapy to alleviate adverse effects of conventional therapy and to improve quality of life (Cheetham 2001; Nam 1999; Petru 2001; Sehouli 2000). This high number of cancer patients using selenium supplements contrasts with the little clear evidence in this field thus justifying the need to update this Cochrane review.

## OBJECTIVES

The aim of this review was to assess the effects of supplementary selenium, given at any dose, on:

- adverse effects of conventional radiotherapy, chemotherapy, and/or oncological surgery;
- quality of life and performance status during and after conventional oncological therapy.

A further aim was to make recommendations for future research.

## METHODS

### Criteria for considering studies for this review

#### Types of studies

Randomised-controlled trials (RCTs) were included in this review which studied the efficacy of selenium supplementation as supplementary cancer treatment. No restriction was made regarding publication status, year or language of publication.

#### Types of participants

Study participants were patients of any age with malignant diseases (regardless of the stage of disease) who were undergoing tumour specific therapy (chemotherapy, radiotherapy or surgery).

#### Types of interventions

Studies that assessed selenium supplements or selenium preparations at any dose, duration, and route of administration compared with placebo or no intervention/routine care were eligible. Studies

using selenium supplements as part of a multicomponent preparation in which there was no study arm testing for selenium supplements alone were not included in this review. Collateral interventions (e.g. manual therapy for lymphoedema) were allowed if adequately controlled for, and used in, all arms of the study.

### Types of outcome measures

Data on the following outcomes were collected:

- the effect of selenium on the incidence and severity of chemotherapy or radiotherapy related toxicities if they were reported according to internationally accepted criteria for common toxicities (e.g. WHO, ECOG or NIH criteria for adverse effects; [DCTD 2004](#); [Miller 1981](#); [Oken 1982](#));
- the effect of selenium on the incidence and severity of lymphoedema following surgical treatment or radiotherapy;
- patient-reported levels of physical and psychological indices of symptom distress (measured using reliable and valid assessment tools);
- quality of life, as measured by a validated instrument;
- incidence and type of adverse effects.

### Search methods for identification of studies

#### Electronic searches

For this update we searched the following databases, sources of grey literature and registers of clinical trials with the reported search strategies:

- the Cochrane Pain, Palliative & Supportive Care Trials Register (July 2007),
- the Cochrane Database of Systematic Reviews (*The Cochrane Library*, Issue 2, 2007),
- the Database of Abstracts of Reviews of Effects (DARE) (*The Cochrane Library*, Issue 2, 2007),
- the Cochrane Central Register of Controlled Trials CENTRAL (*The Cochrane Library*, Issue 2, 2007),
- MEDLINE (1966 to July 2007),
- EMBASE (1980 to 2007 week 28),
- SIGLE (October 2004; database discontinued in 2005),
- Cancerlit (July 2007),
- CCMed (Clinical Contents in Medicine, German) (October 2004),
- the German online register for cancer trials (July 2007),
- the International Standard Randomised Controlled Trial Number Register ISRCTN (October 2004),
- the Meta-Register of Controlled Trials mRCT (October 2004),
- the NCI Clinical Trials register (October 2004).

The search for the original review was run in 2004, subsequent searches for the update were run in 2007. CCMed database - a

database of German journals - search was omitted in 2007 as the relevant journals for this review are now indexed in MEDLINE, which they were not in 2004. ISRCTN, mRCT and NCI Clinical Trials are online registers for clinical studies and helpful to identify ongoing RCTs. We considered a time interval of three years too short for an update search in these online resources taking into account a reasonable balance of resources and expected outcome. We used the MEDLINE search strategy as outlined in [Appendix 1](#) (via PubMed - 1966 to July 2007), its strategy was adapted and developed for other databases searched which can be seen in [Appendix 2](#).

#### Searching other resources

We contacted the Chinese Cochrane Center and the Russian Branch of the Nordic Cochrane Center, but due to limited resources we were not able to have the Moscow Medical Library or the Chinese Biomedical Database searched for publications. Additionally, we checked the reference lists of retrieved publications for further studies and searched the archive of our work group. Manufacturers of selenium supplements and authors of publications were contacted and asked to contribute additional data.

### Data collection and analysis

#### Selection of studies

Identified publications were checked for eligibility by both review authors. A publication was obtained in full text if it could not be rejected with certainty on the basis of its title or abstract. Studies of possible relevance were also obtained in full text for further evaluation. If inclusion or exclusion could not be assessed from the publication alone, first authors were contacted for additional information. Reasons for excluding trials from the review are reported in the '[Characteristics of excluded studies](#)' table. The review authors were not blinded to authors' names, institution and source of publication. Any disagreement was resolved by discussion between the review authors.

#### Data extraction and management

Both review authors independently extracted and documented data using a pre-tested extraction form. The following information was collected:

- study methods (randomisation, allocation concealment, blinding, eligibility criteria, follow-up);
- participants (patient characteristics, age, gender, cancer type, other cancer management/treatment, comorbidity) and additional treatments/medication;

- interventions (selenium dose, regimen, duration of treatment, route of administration) and placebo
- outcome measures;
- study withdrawals, drop-outs and protocol deviations;
- adverse effects;
- informed consent, ethic board approval.

Discrepancies were resolved by discussion between both review authors.

We contacted all primary study authors of included studies for additional information and data. Details of all eligible studies are summarised in the 'Characteristics of included studies' table.

### Methodological quality

Review authors assessed studies for methodological quality using a critical appraisal checklist according to [Juni 2001](#).

This checklist included the following aspects of methodological quality:

- randomisation, allocation concealment,
- blinding,
- specification of eligibility criteria,
- equal provision of care (apart from selenium supplementation) and equal follow-up,
- similarity of groups at baseline,
- reporting of protocol deviations, withdrawals and drop-outs,
- reporting of outcome measures and adverse effects,
- statistical procedures (intention-to-treat analysis).

Additionally, completed studies were rated according to the Oxford Quality Scale ([Jadad 1996](#)) and the Delphi list ([Verhagen 1998](#)).

### Analysis

We planned to calculate a summary statistic and its 95% confidence interval (CI) for each outcome and to perform a meta-analysis of summary statistics if appropriate. Due to the small number of included studies, their methodological limitations and clinical heterogeneity, we considered summary statistics not to be appropriate. Study results are reported as presented in the original publication along with information on adverse effects.

## RESULTS

### Description of studies

#### Results of the search

For this update review one new included study ([Zimmermann 2005](#)) was included and more participants were added to the [Mücke 2007](#) study adding a further 39 participants to the total participants included within the review. The previous review identified an ongoing study which has been completed since the original publication of this review and is now excluded ([Schumacher 2003](#)). The total number of participants considered within the included studies is 162.

Our search returned 1981 studies of possible relevance. Of these, 1943 studies could be excluded based on title or abstract. The apparently high number of screened studies was due to our intensive literature search in databases that did not allow the use of a search strategy other than the main topic "selenium". After screening, the remaining 38 publications were obtained in full text. Following the inclusion criteria of this review, a further 30 papers (representing 26 studies) had to be excluded. The main reasons for exclusion were that the publications described a non-randomised study or the study investigated laboratory parameters, e.g. the change of serum selenium status after supplementation. (see 'Characteristics of excluded studies').

The remaining eight publications represented four studies ([Büntzel 2004](#); [Kasseroller 1998](#); [Mücke 2007](#); [Zimmermann 2005](#)). Two of these studies were completed ([Kasseroller 1998](#); [Zimmermann 2005](#)) and intermediate results have been published for one of the ongoing studies ([Mücke 2007](#)) (see 'Characteristics of included studies'). One study is ongoing ([Büntzel 2004](#)) and no preliminary results have been published so far. All study authors were contacted and provided additional papers or information for this review.

All four studies were or are being conducted in Germany and Austria as a cooperation between different clinicians and one German pharmaceutical company, which manufactures sodium selenite products for clinical application.

### Included studies

#### 1) Effects on the incidence and severity of lymphoedema

Two completed RCTs could be identified for this outcome ([Kasseroller 1998](#); [Zimmermann 2005](#)):

[Kasseroller 1998](#) was described as a placebo-controlled randomised, double blind trial. Sixty female cancer patients were included in the study and results were presented for 57. The objective was to investigate the efficacy of sodium selenite application in combination with physical therapy to relieve secondary lymphoedemas after breast cancer surgery; recurrence of erysipela infection in the oedematous limb was reported as the primary outcome. Participants in the intervention group received sodium selenite per os for 15 weeks with a sum dose of 19.600 µg to 28.000 µg (19.6 mg to 28 mg, depending on body weight). All participants were treated with a combined physical congestion-relief programme.

Preliminary results of this study were published in 1996 giving data for 34 female cancer patients, including 25 women after breast cancer surgery and nine women after pelvic surgery. According to our correspondence with the author, all 25 breast cancer patients reported in 1996 were participants of this later study (Kasseroller 1998). We could not obtain additional information on the remaining participants with other gynecologic malignancies. Zimmermann 2005 was also described as a placebo-controlled randomised, double-blind trial. Twenty participants (18 male, two female) with head-and-neck cancer undergoing curatively intended tumour resection were included. The objective was to investigate blood selenium concentration and different enzyme activities and postoperative lymphoedema. Only the latter outcome is of interest for this systematic review. Participants in the intervention group received 3000 µg sodium selenite intravenously on the day of surgery and 1000 µg sodium selenite daily (i.v. or per os (p.o.)) on days one to 21 after surgery.

## 2) Effects on the incidence and severity of chemo- or radiotherapy related toxicities

One ongoing RCT with published preliminary results could be identified for this outcome:

Mücke 2007 investigated the use of sodium selenite supplements in female cancer patients during adjuvant radiotherapy in an ongoing randomised, unblinded two-armed multi-center study with a no-treatment control group. Study objectives were the assessment of the efficacy of sodium selenite p.o. for normalization of selenium deficiency during adjuvant radiotherapy (primary outcome) and, secondarily, its effect on the incidence and severity of radiotherapy toxicities (e.g. diarrhoea according to common toxicity criteria (CTC), changes in body weight and blood count). The intervention group received a sum dosage of approximately 15,900 to 18,000 µg sodium selenite p.o. (15.9 to 18 mg) during radiotherapy (ca. 5 weeks). According to the study protocol, the trial will randomise 200 participants to intervention and control group. Intermediate results have been published for 41 participants (Mücke 2004b) and 63 participants (Mücke 2004a), 77 participants (Mücke 2006b), 79 participants (Mücke 2006a), 80 participants (Mücke 2007), and 82 participants (Mücke 2007a; Mücke 2007b) (Mücke 2007).

## Risk of bias in included studies

Details of the methodological quality of included studies are given in an additional table (see Additional Table 1: 'Quality Assessment of Included Studies').

No study described the method of sequence generation for random allocation. Mücke 2007 reported an adequate procedure of allocation concealment (personal communication); methods of allocation concealment in Kasseroller 1998 and Zimmermann 2005 remained unclear. All studies specified criteria for inclusion and

exclusion of participants, but information on intervention and control groups' characteristics at baseline was not provided. Therefore, unequal distribution of risk factors for the outcomes of interests cannot be ruled out. Treatment allocation was described as blinded to participants, doctors, care-givers and outcome assessors in Kasseroller 1998, but a control procedure checker as to whether blinding was successful was not reported. Zimmermann 2005 was described as a double-blinded, placebo-controlled trial. However, the informed consent form for participants indicated that they had been allocated to the selenium group (see: Leonhardt 2002, p. 79 (Zimmermann 2005)). We were not able to clarify the actual procedure of treatment allocation and blinding with the investigators of this study. Mücke 2007 is an unblinded trial.

According to Kasseroller 1998, all participants received comparable physical treatment for lymphoedema. However, details on concomitant treatments such as high-voltage therapy and information on additional medication that might influence erysipela incidence (e.g. corticosteroids) were not provided. All participants in Zimmermann 2005 underwent curatively intended oral tumour surgery with bilateral neck dissection. Information on factors that might influence postsurgical local swelling, such as the extent of intraoperative tissue trauma, postoperative complications or comorbidity was not reported. In the ongoing trial by Mücke 2007, participants are excluded who receive specific medication at study commencement (antibiotics, antacids, adsorbents, 5-HT<sub>3</sub>-antagonists). If started after study inclusion, concomitant intake of these drugs should be avoided according to the study protocol, but will not result in exclusion from the study group. Other drugs that might influence diarrhoea, especially opioid analgesics, are not mentioned in the study protocol and not reported in the preliminary data.

Kasseroller 1998 gave no details for methods of outcome assessment or criteria of erysipela diagnosis. Especially the assessment of erysipela infections after dismissal from hospital remained unclear. Zimmermann 2005 reported three distances in the face (tragus - nostril, tragus - tip of chin, tragus - corner of mouth), each distance measured both directly and circumferentially. The measurements were manually undertaken by one investigator before surgery and repeated immediately after, one and two weeks after tumour resection. Reliability of measurements was unclear. Study investigators equated the measurement of postsurgical face swelling within the first two weeks after surgery with the assessment of postoperative secondary lymphoedema. The validity of this procedure seems questionable as the short interval between surgery and outcome assessment does not allow for postsurgical traumatic swelling to disappear and for lymphoedema to develop. In addition, Zimmermann and colleagues themselves stated that the "measuring distances chosen only partly describe the extent of the lymphoedema" (Zimmermann 2005, p. 199). Measurements of caudal to mandible, where lymphoedema would be most prominent according to Zimmermann and colleagues, were not conducted. In Mücke 2007, incidence and severity of diarrhoea are

diagnosed by the radiologist on a weekly basis based on participants' self-reports.

[Kasseroller 1998](#) reported findings for recurrence of erysipela infection. In contrast to the study objectives, results on the development of lymphoedema with and without selenium supplementation were not reported. Results of a data analysis (i.e. OR/CI or t-test/P) were not presented. [Zimmermann 2005](#) reported the measurements for three distances in the face at four time points as raw data. Results were presented as graphs only; Cross-tables or absolute numbers of results were not provided. Results were tested for differences between intervention and control group and significance levels were reported; the related test was not specified. [Mücke 2007](#) reported preliminary data on incidence and severity of radiation-induced diarrhoea and chi-square test and t-test were performed to investigate differences in the occurrence of diarrhoea according to CTC.

## Effects of interventions

After a comprehensive literature search, we located two completed studies investigating the effect of selenium supplementation on secondary lymphoedema. Another ongoing study examines radiotherapy-associated diarrhoea as a secondary outcome; with the primary objective being the evaluation of sodium selenite supplementation for compensation of pre-existing selenium deficiency. We could not identify trials covering other outcomes of this review, such as quality of life or chemotherapy toxicities.

### 1) Effects of selenium with physical therapy on the incidence and severity of lymphoedema

[Kasseroller 1998](#) measured the recurrence of erysipela infections of lymphoedematous upper limbs after breast cancer treatment. During the three-week treatment in the hospital no erysipela occurred in the intervention group, and one erysipela occurred in the control group. During the entire intervention and follow-up period of 15 weeks no participant in the intervention group developed recurrent erysipela while in the placebo group 14 participants (50 % of n = 28) were diagnosed with erysipela infections. These findings suggest that there might be a lower incidence of recurrent erysipela infections in the selenium group compared to placebo in the 15 weeks' period. However, this study has severe methodological limitations, as described earlier in this review. Due to these methodological problems, no conclusions on the effect of selenium supplementation on the recurrence of erysipelas in the intervention group can be drawn and generalisations to other cancer patients seem questionable.

[Zimmermann 2005](#) reported the extent of facial swelling after oral tumour surgery for head and neck cancer. The control group showed the peak in facial swelling at the measuring point one week after surgery. In the selenium group, the maximum swelling

was seen directly after surgery. The circumferential distance tragus/tip of chin was significantly shorter one week and two weeks after surgery in the intervention group, indicating a lesser facial swelling than in the control group. Taking into consideration the low number of participants and the limitations in methodology and reporting, both internal validity of results and external generalisability to other patients seems questionable.

### 2) Effects of selenium on the incidence and severity of chemo- or radiotherapy related toxicities

[Mücke 2007](#) investigated radiotherapy associated diarrhoea as a secondary outcome in selenium deficient women undergoing pelvic radiation. The latest publication of this study provided data on 82 female participants with a median age of 66 years (range 31 to 80). Authors reported a statistically significant lower incidence and severity of diarrhoea in weeks four to six of radiotherapy in the intervention group. The overall incidence of radiation-induced diarrhoea grade CTC two or higher was 21% in the selenium group compared to 47 % in the control group. Further data on incidence or severity of diarrhoea, cross-tables, additional data analysis (e.g. OR with CI) or results (e.g. blood count) have not yet been presented. Also, distribution of risk factors for diarrhoea at baseline and concomitant medication that might influence diarrhoea have not been reported. A conclusive discussion of this trial will need to await the publication of detailed final results.

In all three studies, no adverse effects of selenium supplementation were observed.

## DISCUSSION

After a comprehensive literature search, we were able to locate only two completed RCTs ([Kasseroller 1998](#); [Zimmermann 2005](#)) that fulfilled the inclusion criteria of this review and another ongoing study ([Mücke 2007](#)) with published preliminary results. All three included studies used inorganic sodium selenite as medication. Another study not included within the included studies is currently ongoing ([Büntzel 2004](#)).

Due to methodological limitations it is not possible to draw conclusions on the effect of selenium supplementation on the recurrence of erysipela infections in women with secondary lymphoedema in the [Kasseroller 1998](#) study. This study has been included in another Cochrane Review assessing the efficacy of anti-inflammatories for reducing acute inflammatory episodes in lymphoedema of the limbs ([Badger 2004](#)), which came to a similar conclusion regarding trial quality and the subsequent limitations in the interpretation of trial results. However, [Badger 2004](#) reported two publications of [Kasseroller 1996](#) and [Kasseroller 1998](#) as two distinct studies. Having been able to make contact with

Kasseroller, it was confirmed that the publication Kasseroller 1996 reported preliminary results of [Kasseroller 1998](#).

[Zimmermann 2005](#) reported a lesser facial swelling after head and neck tumour resection in one (of three) measurements in a two week follow-up in a group of 20 participants. Considering the unclear reliability and validity of outcome measurements, it is unclear whether the measured differences really reflect a reduction in postoperative lymphoedema in the selenium group. Generalisation to other cancer patients seems questionable. Furthermore, the clinical relevance of these measurements remained unclear.

The findings of Kasseroller have been cited in a number of secondary publications and non-systematic reviews. In 2000 Kasseroller/Schrauzer published an overview with pooled results from different studies that investigated the efficacy of selenium supplementation in female breast cancer patients with secondary lymphoedema ([Kasseroller 2000](#)). We obtained confirmation from Kasseroller that his study data ([Kasseroller 1998](#)) were included in this summary report, but could not clarify where the remaining data stemmed from. It is doubtful that all data in this overview originated from RCTs. Kasseroller/Schrauzer inferred that clinical “studies with lymphedema of the arm after breast cancer surgery document that sodium selenite increases the efficacy of physical decongestion therapy and prevents erysipela infections” ([Kasseroller 2000](#)). Another summary of current treatment options for secondary lymphoedema ([Bruns 2003](#)) arrived at a similar positive conclusion regarding selenium supplementation: “Present data demonstrate that selenium can enhance the benefits of physical therapy in radiation-induced lymphedemas. The very low toxicity profile of selenium and its cost effectiveness are further arguments for its use in lymphedema treatment” ([Bruns 2003](#)). Authors backed their approval with a reference to the [Kasseroller 1998](#) study and asserted that “Kasseroller [...] reported promising results of a placebo-controlled, double-blind study of selenium in 179 postmastectomy patients”. However, the [Kasseroller 1998](#) study included only 60 participants and the above described overview by Kasseroller/Schrauzer ([Kasseroller 2000](#)), which reported data of 179 participants altogether, did not describe a RCT, but pooled data of predominantly unknown origin.

More recent publications also referred to [Zimmermann 2005](#) to back their approval of selenium supplementation ([Beuth 2007](#)) claiming that “[r]andomized controlled clinical trials have demonstrated significant benefits for cancer patients receiving Se during chemo and radiotherapy, such as reduction of lymphedema in head and neck” cancer patients (p. 427).

Considering the lack of evidence from studies with a design that aimed to minimise possible sources of bias and the low quality of reporting in the two completed trials, this systematic review does not support the conclusions in the above cited secondary publications on the efficacy of selenium supplementation in patients with secondary lymphoedema.

To date no completed RCT is available that investigates the efficacy of selenium supplements to ameliorate side effects of conventional cancer chemo- or radiotherapy. Intermediate results from one study using sodium selenite to diminish diarrhoea in women undergoing pelvic radiation were included in this review, but have not yet been published in sufficient detail for a conclusive discussion ([Mücke 2007](#)). Another study is currently ongoing with toxicities of conventional therapy as outcomes of interest: [Büntzel 2004](#) has a design comparable to that of [Mücke 2007](#) and includes participants with a pre-existing selenium deficiency who are undergoing radiotherapy for ENT-malignancies. This study has been recruiting since 2000 and aims to include 200 participants randomly allocated to intervention (oral administration of sodium selenite) or control group (without selenium supplementation). Primary outcome measure is the normalisation of selenium deficiency in the blood, and secondary outcomes include radiotherapy toxicities.

In contrast to the complete lack of evidence from RCTs regarding the efficacy of selenium supplements against radio-/chemotherapy associated side-effects, selenium supplements are frequently recommended for this indication in secondary publications ([Arnold 2001](#); [Beuth 2002](#)) and claimed that “a protective effect against chemotherapy toxicities” has been proven in clinical studies and observations ([Sill-Steffens 2003](#)). Considering the lack of evidence, no recommendation can currently be given regarding the use of selenium supplements to ameliorate radio- or chemotherapy associated side effects.

Authors of secondary publications frequently emphasize that selenium supplementation seems to be a safe intervention as no side effects have been reported in clinical studies. In both RCTs that are included in this review no side effects attributable to selenium supplementation were reported at average dosages between 100 to 1000 µg sodium selenite daily over a period of five to 15 weeks. Inorganic selenium compounds have known acute and chronic toxicities (selenosis) at high dosages. The recommended daily allowance for adults varies between 30 to 100 µg daily depending on country and gender. 200 µg to 300 µg are considered the maximally safe daily intake over longer periods ([Baehr 1999](#)), but dosages of 1000 µg daily have been used in clinical investigations with sepsis patients for up to four weeks without signs of selenosis ([Zimmermann 1997](#)).

A few cases of acute and chronic selenosis in patients due to accidental overdosage of selenium-containing medication have been reported in the literature. In the year 2000 two severe incidents occurred in Germany and Austria when physicians mistakenly prescribed milligrams of sodium selenite instead of micrograms causing one death in Austria after parenteral administration of about 200 mg selenium ([Pfeffer 2002](#)). In 2006, the case of an Australian man with an adenocarcinoma of the prostate was published, who ingested 10 g of sodium selenite after reading on the internet about selenium as putative treatment for prostate cancer. He died some

hours later of selenium intoxication (See 2006). Several cases of chronic selenosis have been reported in persons using non-prescribed nutritional supplements with inaccurate specification of selenium content. Individuals had a daily intake of ca. 1000 µg/d selenium up to 27,000 µg/d in the form of organic or inorganic selenium compounds over several weeks resulting in chronic selenium intoxication with partly severe symptoms (nausea, vomiting, loss of nails and hair, and halitosis) (Helzlsouer 1985; Jensen 1984).

Apart from acute and chronic toxicity of selenium, long-term effects of selenium supplementation are still controversially discussed (Vinceti 2001). Evidence from observational and intervention trials regarding the effect of selenium on cancer incidence and mortality is contradictory; a Cochrane review on the cancer-preventive efficacy of selenium is in process (Dennert 2005). In Germany, inorganic selenium compounds are currently classified as potential workplace carcinogens (Greim 1999; Greim 2003). Whether short or long-term selenium supplementation bears any protective or harmful effect on cancer diseases, is still open to question.

To summarise, the current evidence is insufficient to provide guidelines for clinical practice. The lack of primary study data contrasts remarkably with the amount of secondary publications and the proportion of cancer patients using selenium supplements, especially in German-speaking countries. It remains to be seen whether the ongoing studies will be able to add substantially to the current body of evidence.

## AUTHORS' CONCLUSIONS

### References to studies included in this review

#### Kasseroller 1998 {published and unpublished data}

Kasseroller R. Prophylaxis of erysipelas in secondary lymphedemas with selenium [Erysipelprophylaxe beim sekundären Lymphodem mit Selen]. *Der Allgemeinarzt* 1996;**18**(3):244–7.

\* Kasseroller R. Sodium selenite as prophylaxis against erysipelas in secondary lymphedema. *Anticancer Research* 1998;**18**(3C):2227–30.

#### Mücke 2007 {published and unpublished data}

Micke 2003a: Micke O, Riesenbeck D, Büntzel J. Relevance of selenium supplementation in postoperative radiotherapy of head-neck-tumours and gynecologic pelvic tumours - A multi-center phase III study [Stellenwert der Selen-

## Implications for practice

The currently available evidence is insufficient to assess reliably the possible role of selenium supplementation on toxicities related to oncology treatments, on the development and severity of secondary lymphoedema and on quality of life during antineoplastic treatments. The quality of the studies prohibits us from extrapolating any effects observed to the general population level. Currently, research findings do not provide a basis for any recommendation in favour or against selenium supplementation in cancer patients. Nevertheless, the potential hazards of supplementing a trace mineral should always be kept in mind.

Since the last version of this review the additional participants have not provided additional information to change the conclusions.

## Implications for research

As it is likely that a considerable number of cancer patients will seek “complementary” treatments to alleviate adverse effects connected with conventional therapy, further research is desirable.

Supplementation with sodium selenite is currently under investigation in RCTs and robust and sound replication of the existing trial might be useful. However, no systematic dosage-finding study has been reported so far. An adequate dosage-finding study seems as desirable as adequate reporting of completed, ongoing and future trials.

## ACKNOWLEDGEMENTS

We thank all authors, clinicians and investigators who provided us with additional information and publications for this review. The AG Biologische Krebs therapie is funded by the German Cancer Aid.

## REFERENCES

Substitution bei der postoperativen Strahlentherapie bei Kopf–Hals–Tumoren und gynäkologischen Beckentumoren – Eine multizentrische Phase–III–Studie]. [www.uni-muenster.de/rektorat/forschungsberichte-2001-2002/fo05aka15.htm](http://www.uni-muenster.de/rektorat/forschungsberichte-2001-2002/fo05aka15.htm) (download 09.01.2004). 48149 Münster: Westfälische Wilhelms Universität, 2003.

Micke 2007: Micke O, Mücke R, Bruns F, Kisters K, Büntzel H. Some clinical results on selenium in radiation oncology. Letter by O Micke, R Mücke, F Bruns, K Kisters, J Büntzel on W. Dörr: Effects of selenium on radiation responses of tumor cells and tissue. *Strahlentherapie und Onkologie* 2007;**183**(6):344–5.

Mücke 2004a: Mücke R, Glatzel M, Bernd-Skorka R, Seifert G, Büntzel J, Schönekaes KG, et al. Adjuvant treatment with sodium selenite in gynecologic radiation

oncology - first results of a phase III study. Abstract for the ESTRO meeting 2004 in Amsterdam. 2004.

Mücke 2004b: Mücke R, Glatzel M, Riesenbeck D, Bernd-Skorka R, Seifert G, Büntzel J, et al. Sodium selenite in gynecologic radiation oncology - first results of a phase III study. *Trace Elements and Electrolytes* 2004;**21**(2):78–82.

Mücke 2004c: Mücke R. Observation plan for the post-marketing surveillance stud: Use of selenase as complementary therapy to postoperative adjuvant radiotherapy of cervical and endometrial carcinoma. Version 1.4 [Beobachtungsplan der Anwendungsbeobachtung (AWB): Einsatz von selenase als Zusatztherapie zur postoperativen adjuvanten Radiotherapie bei Zervix- und Endometriumkarzinom. Version 1.4]. Biosyn Arzneimittel GmbH 2004.

Mücke 2004d: Mücke R. Use of selenase as supplementary therapy in postoperative adjuvant radiotherapy for cervical and endometrial carcinoma [Einsatz von selenase als Zusatztherapie zur postoperativen adjuvanten Radiotherapie bei Zervix- und Endometriumkarzinom]. <http://www.studien.de/>. 92637 Weiden, Söllnerstr.16: Dr. Ralph Mücke, Klinikum Weiden, download 16.01.2004.

Mücke 2006a: Mücke R, Glatzel M, Reichl B, Bernd-Skorka R, Baaske D, Kaiser A, et al. V 6 Sodium selenite in gynecologic radiation oncology - results of a prospective randomized observation study. AKTE. Münster: Diplodocus, 2006; Vol. Band 8.

Mücke 2006b: Mücke R, Glatzel M, Reichl B, Bernd-Skorka R, Büntzel J, Bruns F, et al. Sodium selenite in gynecologic radiation oncology: Results of a prospective randomized observation study. *Trace Elements and Electrolytes* 2006;**23**(3):173–7.

\* Mücke 2007a: Mücke R. Impact of Selenium in Radiation Oncology. Conference presentation at the 23rd Annual Meeting of the Society for Mineral and Trace Elements, 28.-29.09.2007 in Stuttgart/Germany 2007.

Mücke 2007b: Mücke R. The impact of sodium selenite in gynecologic radiation oncology - final results of a prospective randomized observation study. Submission abstract for conference presentation at the 23rd Annual Meeting of the Society for Mineral and Trace Elements 28.-29.09.2007 in Stuttgart/Germany 2007.

#### **Zimmermann 2005 {published and unpublished data}**

Leonhardt 2002: Leonhardt H. Impact of sodium selenite on the postoperative chronic lymphoedema and inflammatory cytokines in patients with stage T3 and T4 carcinomas of the base of the mouth and tongue [Wirkung von natrimselenit auf das postoperative chronische lymphödem und inflammatorische zytokine bei patienten mit mundboden- und zungengrundkarzinomen der stadien T3 und T4]. Thesis. Dresden: Technischen Universität Dresden, 2002.

\* Zimmermann T, Leonhardt H, Kersting S, Albrecht S, Range U, Eckelt U. Reduction of postoperative lymphedema after oral tumor surgery with sodium selenite. *Biological Trace Element Research* 2005;**106**(3):193–203.

## References to studies excluded from this review

#### **Asfour 2006 {published data only}**

Asfour IA, El SS, Fayek MH, Hegab HM, Raouf S, Moussa MA. Effect of high-dose sodium selenite therapy on polymorphonuclear leukocyte apoptosis in non-Hodgkin's lymphoma patients. *Biological Trace Element Research* 2006;**110**(1):19–32.

#### **Badger 2004 {published data only}**

Badger C, Seers K, Preston N, Mortimer P. Antibiotics/anti-inflammatories for reducing acute inflammatory episodes in lymphoedema of the limbs. *Cochrane Database of Systematic Reviews* 2004, Issue 2. [DOI: 10.1002/14651858.CD003143.pub2

#### **Büntzel 1999 {published data only}**

Büntzel J. Experiences with sodium selenite in treatment of acute and late adverse effects of radiochemotherapy of head-neck carcinomas. Cytoprotection working group in AK supportive measures in oncology within the scope of MASCC and DKG [Erfahrungen mit Natriumselenit in der Behandlung von akuten und späten Nebenwirkungen der Radiochemotherapie von Kopf-Hals-Karzinomen.]. *Medizinische Klinik* 1999;**94**(Suppl 3):49–53.

#### **Elango 2006 {published data only}**

Elango N, Samuel S, Chinnakkannu P. Enzymatic and non-enzymatic antioxidant status in stage (III) human oral squamous cell carcinoma and treated with radical radiotherapy: influence of selenium supplementation. *Clinica Chimica Acta* 2006;**373**(1-2):92–8.

#### **Fakih 2006 {published data only}**

Fakih MG, Pendyala L, Smith PF, Creaven PJ, Reid ME, Badmaev V, et al. A phase I and pharmacokinetic study of fixed-dose selenomethionine and irinotecan in solid tumors. *Clinical Cancer Research* 2006;**12**(4):1237–44.

#### **Federico 2001 {published data only}**

Federico A, Iodice P, Federico P, Del Rio A, Mellone MC, Catalano G, et al. Effects of selenium and zinc supplementation on nutritional status in patients with cancer of digestive tract. *European Journal of Clinical Nutrition* 2001;**55**(4):293–7.

#### **Fraunholz 2001 {published data only}**

Fraunholz I, Jüling-Pohlitz L, Böttcher H. Influence of selenium on acute mucositis in radiochemotherapy of head and neck tumors. *Trace Elements and Electrolytes YR*: 2001. 18 Vol. 18:98–9.

#### **Hoenjet 2005 {published data only}**

Hoenjet KM, Dagnelie PC, Delaere KP, Wijckmans NE, Zambon JV, Oosterhof GO. Effect of a nutritional supplement containing vitamin E, selenium, vitamin c and coenzyme Q10 on serum PSA in patients with hormonally untreated carcinoma of the prostate: a randomised placebo-controlled study. *European Urology* 2005;**47**(4):433–9.

**Hu 1997** *{published data only}*

Hu Y, Lin Y, Xuemei S. Clinical study on the effect of kappa-selenocarrageenan in reducing the nephrotoxicity of cis-platin. *Chinese New Drug Journal* 1997;**6**:349–51.

\* Hu YJ, Chen Y, Zhang YQ, Zhou MZ, Song XM, Zhang BZ, et al. The protective role of selenium on the toxicity of cisplatin-contained chemotherapy regimen in cancer patients. *Biological Trace Element Research* 1997;**56**:331–41.

**Kasseroller 1995** *{published data only}*

Kasseroller R. Sodium selenite in the therapy of chronic lymphedema [Natriumselenit in der Therapie des chronischen Lymphödems]. *Der Allgemeinarzt* 1995;**17** (13):1396–404.

**Kiremidjian 2000** *{published data only}*

Kiremidjian-Schumacher L, Roy M. Effect of selenium on the immunocompetence of patients with head and neck cancer and on adoptive immunotherapy of early and established lesions. *BioFactors* 2001;**14**(1-4):161–8.

\* Kiremidjian-Schumacher L, Roy M, Glickman R, Schneider K, Rothstein S, Cooper J, et al. Selenium and immunocompetence in patients with head and neck cancer. *Biological Trace Element Research* 2000;**73**(2):97–111.

**Kreienberg 2003** *{published data only}*

Kreienberg R. Prospective, randomised, 4-armed, controlled, open, multi-center phase-II-study: influence of selenium on the adverse effect profile of adjuvant chemotherapy in patients with mammary carcinoma [Prospektive, randomisierte, vierarmige, kontrollierte, offene, multizentrische Phase-II-Prüfung: Einfluss von Selen auf das Nebenwirkungsprofil von adjuvanter Chemotherapie bei Patientinnen mit Mamakarzinom]. online study register University of Ulm, Department for Gynecology: www.uni-ulm.de 2003.

**Lasch 1999** *{published data only}*

\* Lasch K, Bräsel C, Jahn H. Selenium therapy in colorectal tumors? [Selentherapie bei kolorektalen Tumoren?]. *Medizinische Klinik* 1999;**94**(Suppl 3):97–100.

Lasch K, Bräsel C, Jahn H. Selenium therapy for colorectal carcinoma [Poster: Selentherapie bei kolorektalen Karzinomen]. Poster. —.

**Schumacher 2003** *{published data only}*

Schumacher K. Influence of selenium on the adverse effect profile of adjuvant chemotherapy in patients with mammary carcinoma. Prospective, controlled, open, multi-center clinical study phase II [Einfluss von Selen auf das Nebenwirkungsprofil von adjuvanter Chemotherapie bei Patientinnen mit Mamakarzinom. Prospektive, kontrollierte, offene, multizentrische klinische Prüfung der Phase II]. www.studien.de (download 16.01.2004) 2002.

\* Schumacher K, Biosyn Arzneimittel GmbH. Study plan - Influence of selenium on the adverse effect profile of adjuvant chemotherapy in patients with mammary carcinoma. Prospective, randomised, 4-armed, controlled, open, multi-center clinical study phase II [Prüfplan – Einfluss von Selen auf das Nebenwirkungsprofil von adjuvanter Chemotherapie bei Patientinnen mit Mamakarzinom.

Prospektive, randomisierte, vierarmige, kontrollierte, offene, multizentrische klinische Prüfung der Phase II]. biosyn Arzneimittel GmbH 2003.

**Sieja 2000** *{published data only}*

Sieja K. Protective role of selenium against the toxicity of multi-drug chemotherapy in patients with ovarian cancer. *Pharmazie* 2000;**55**(12):958–9.

**Sundström 1984** *{published data only}*

Sundström H, Korpela H, Viinikka L, Kauppila A. Serum selenium and glutathione peroxidase, and plasma lipid peroxides in uterine, ovarian or vulvar cancer, and their responses to antioxidants in patients with ovarian cancer. *Cancer Letters* 1984;**24**(1):1–10.

**Sundström 1989** *{published data only}*

Sundström H, Korpela H, Sajanti E, Kauppila A. Supplementation with selenium, vitamin E and their combination in gynaecological cancer during cytotoxic chemotherapy. *Carcinogenesis* 1989;**10**(2):273–8.

**Wagler 2000** *{published data only}*

Wagler E, Schwokowski C, Schönfelder M. Poster: first results of selenium supplementation in patients with gastrointestinal tumours [Poster: Erste Ergebnisse der Selensubstitution bei Patienten mit gastrointestinalen Tumoren]. Poster. Universitätsklinikum Leipzig, 2000.

**Xu 1990** *{published data only}*

Xu H, Mei W, Dong Z, Liao B. Study of the oxidative metabolic function and chemotaxis of neutrophils from patients with cancer influenced by selenium yeast. *Biological Trace Element Research* 1990;**25**(3):201–9.

**Xu 1999** *{published data only}*

Xu B, Sun Y. Randomized double-blind placebo-controlled phase II clinical trial of kappa-selenocarrageenan in cancer chemotherapy patients. *Chinese Journal of Clinical Oncology* 1999;**26**(9):678–80.

**Yu 1996** *{published data only}*

Yu B, Wang M, Li D. [The relationship between selenium and immunity in large bowel cancer]. *Zhonghua Wai Ke Za Zhi* 1996;**34**(1):50–3.

**References to ongoing studies****Büntzel 2004** *{published data only}*

Büntzel 2002: Büntzel J. Use of selenase as complementary treatment to radiotherapy for tumours of the head and neck region [Einsatz von selenase als Zusatztherapie zur Radiotherapie bei Tumoren im Hals–Nasen–Ohrenbereich]. www.studien.de (retrieved on 16.01.2004) 2002.

\* Büntzel J, Biosyn Arzneimittel GmbH. Observation plan of a post-marketing surveillance study. Use of selenase as complementary treatment to radiotherapy of tumours of the head and neck region (version 1.4) [Beobachtungsplan der Anwendungsbeobachtung (AWB). Einsatz von selenase als Zusatztherapie zur Radiotherapie bei Tumoren im Hals–Nasen–Ohrenbereich (version 1.4.)]. biosyn Arzneimittel GmbH 2004.

Micke 2003b: Micke O, Riesenbeck D, Büntzel J.

Importance of selenium supplementation in postoperative

radiotherapy of head-neck-tumours and gynecological tumours of the pelvis - A multi-centre phase III study [Stellenwert der Selen-Substitution bei der postoperativen Strahlentherapie bei Kopf-Hals-Tumoren und gynäkologischen Beckentumoren – Eine multizentrische Phase-III-Studie]. [www.uni-muenster.de/~rektorat/forschungsberichte-2001-2002/fo05aka15.htm](http://www.uni-muenster.de/~rektorat/forschungsberichte-2001-2002/fo05aka15.htm) (download 09.01.2004). Münster: Westfälische Wilhelms Universität, 2003.

## Additional references

### Arnold 2001

Arnold-von Versen B. Use of micronutrients for tumour diseases and tumour prevention [Der einatz von mikronährstoffen bei tumorerkkrankungen und zur tumorprävention]. *Deutsche Zeitschrift für Onkologie* 2001; **33**:19–29.

### Baehr 1999

Baehr K, Dreher I, Koehrl J. Selenium supplementation by selenium yeast and sodium selenite: Analysis of the selenium status as well as risks of deficiency and intoxication [Selensupplementation durch selenhefe und natriumselenit: analyse des selenstatus sowie risiken des mangels und der intoxication]. *The Journal of Laboratory and Clinical Medicine* 1999; **23**(11):594–9.

### Baldew 1989

Baldew GS, van den Hamer CJ, Los G, Vermeulen NP, De Goeij JJ, McVie JG. Selenium-induced protection against cis-diamminedichloroplatinum (II) nephrotoxicity in mice and rats. *Cancer Research* 1989; **49**(11):3020–3.

### Beuth 2002

Beuth J, Uhlenbruck G. Evidence-based complementary therapy measures for prostate carcinoma [Evidenzbasierte komplementäre therapiemaßnahmen beim prostatakarcinom]. *Deutsche Zeitschrift für Onkologie* 2002; **34**:86–90.

### Beuth 2007

Beuth J, Schierholz JM. Evidence-based complementary oncology. Innovative approaches to optimize standard therapy strategies. *In Vivo* 2007; **21**(2):423–8.

### Boucher 1995

Boucher F, Coudray C, Tirard V, Barandier C, Tresallet N, Favier A, de Leiris J. Oral selenium supplementation in rats reduces cardiac toxicity of adriamycin during ischemia and reperfusion. *Nutrition* 1995; **11**(5 Suppl):708–11.

### Bruns 2003

Bruns F, Micke O, Bremer M. Current status of selenium and other treatments for secondary lymphedema. *The Journal of Supportive Oncology* 2003; **1**(2):121–30.

### Cheetham 2001

Cheetham PJ, Le Monnier KJ, Brewster SF. Attitudes and use of alternative therapies in UK prostate cancer patients - isn't it time we were in the know?. *Prostate Cancer and Prostatic Diseases* 2001; **4**:235–41.

### DCTD 2004

DCTD, NCI, NIH, DHHS. Cancer therapy evaluation program: Common terminology criteria for adverse effects. <http://ctep.cancer.gov/reporting/ctcnew.html> 2004.

### Dennert 2005

Dennert G, Zwahlen M, Brinkmann M, Vinceti M, Zeegers MPA, Horneber MA. Selenium for preventing cancer. *Cochrane Database of Systematic Reviews* 2005, Issue 2. [DOI: 10.1002/14651858.CD005195]

### Dietz 1998

Dietz A, Rudat V, Nollert J, Helbig M, Vanselow B, Weidauer H. The chronic larynx oedema as delayed reaction after radiochemotherapy [Das chronische larynxödem als spätreaktion nach radiochemotherapie]. *HNO* 1998; **46**:731–8.

### Erickson 2001

Erickson VS, Pearson ML, Ganz PA, Adams J, Kahn KL. Arm edema in breast cancer patients. *Journal of the National Cancer Institute* 2001; **93**(2):96–111.

### Francescato 2001

Francescato HD, Costa RS, Rodrigues Camargo SM, Zanetti MA, Lavrador MA, et al. Effect of oral selenium administration on cisplatin-induced nephrotoxicity in rats. *Pharmacological Research* 2001; **43**(1):77–82.

### Greim 1999

Greim H. Selenium and its anorganic compounds [Selen und seine anorganischen Verbindungen]. *Gesundheitsschädliche Arbeitsstoffe. Toxikologisch-arbeitsmedizinische Begründungen von MAK-Werten*. Vol. **Looseleaf: Supplement**, Weinheim: Wiley-VCH, 1999.

### Greim 2003

Greim H. Selenium and its inorganic compounds [Selen und sein anorganischen Verbindungen]. In: Greim H editor(s). *Gesundheitsschädliche Arbeitsstoffe. Toxikologisch-arbeitsmedizinische Begründungen von MAK-Werten*. Vol. **Looseleaf: Supplement**, Weinheim: Wiley-VCH, 1999–2003.

### Groetz 2003

Groetz KA, Riesenbeck D, Doerr W. The oral cavity of patients with head-neck-radiotherapy - III. Postradiotherapeutic care [Die mundhöhle des patienten mit kopf-hals-strahlentherapie – III. Postradiotherapeutische betreuung]. *Im Focus Onkologie* 2003; **7-8**:58–61.

### Hatfield 2001

Hatfield DL (Ed). *Selenium. Its molecular biology and role in human health*. Boston/Dordrecht/London: Kluwer Academic Publishers, 2001.

### Helzlsouer 1985

Helzlsouer K, Jacobs R, Morris S. Acute selenium intoxication in the United States. *Federation Proceedings* 1985; **44**:1670–2.

### Jadad 1996

Jadad AR, Moore RA, Carroll D, Jenkinson C, Reynolds DJM, Gavaghan DJ, et al. Assessing the quality of reports of

- randomized clinical trials: Is blinding necessary?. *Controlled Clinical Trials* 1996;**17**:1–12.
- Jensen 1984**  
Jensen R, Claoson W. Selenium intoxication. *New York Morbidity and Mortality Weekly Report* 1984;**33**:157–60.
- Juni 2001**  
Juni P, Altman DG, Egger M. Assessing the methodological quality of randomised controlled trials. In: Egger M, Smith GD, Altman DG editor(s). *Systematic Reviews in Health Care*. 2nd Edition. London: BMJ Publishing Group, 2001: 87–109.
- Kasseroller 2000**  
Kasseroller RG, Schrauzer GN. Treatment of secondary lymphedema of the arm with physical decongestive therapy and sodium selenite: a review. *American Journal of Therapeutics* 2000;**7**(4):273–9.
- Kotamraju 2000**  
Kotamraju S, Konorev EA, Joseph J, et al. Doxorubicin-induced apoptosis in endothelial cells and cardiomyocytes is ameliorated by nitron spin traps and ebsele: role of reactive oxygen and nitrogen species. *Journal of Biological Chemistry* 2000;**275**:33585–92.
- Micke 2003**  
Micke O, Bruns F, Mucke R, Schafer U, Glatzel M, DeVries AF, et al. Selenium in the treatment of radiation-associated secondary lymphedema. *International Journal of Radiation Oncology Biology Physics* 2003;**56**(1):40–9.
- Miller 1981**  
Miller AB, Hoogstraten B, Staquet M, Winkler A. Reporting results of cancer treatment. *Cancer* 1981;**47**:207–14.
- Nam 1999**  
Nam RK, Fleshner N, et al. Prevalence and patterns of the use of complementary therapies among prostate cancer patients: An epidemiologic analysis. *Journal of Urology* 1999;**161**:1521–4.
- Ohkawa 1988**  
Ohkawa K, Tsukada Y, Dohzono H, Koike K, Terashima Y. The effects of co-administration of selenium and cisplatin (CDDP) on CDDP-induced toxicity and antitumour activity. *British Journal of Cancer* 1988;**58**(1):38–41.
- Oken 1982**  
Oken MM, Creech RH, Tormey DC, Horton J, Davis TE, McFadden ET, Carbone PP. Toxicity and response criteria of the eastern cooperative oncology group. *American Journal of Clinical Oncology* 1982;**5**:649–55.
- Petru 2001**  
Petru E, Schmied P, Petru C. Complementary measures with gynecological cancer patients during chemo- and radiotherapy - review and critical consideration of practical issues [Komplementäre maßnahmen bei patientinnen mit gynäkologischen malignomen unter chemo- und strahlentherapie – bestandsaufnahme und kritische überlegungen für die praxis]. *Geburtshilfe Frauenheilkunde* 2001;**61**:75–8.
- Pfeffer 2002**  
Pfeffer H. Severe incident with selenium medication [Schwerwiegender zwischenfall bei selen-medikation]. *Hamburger Zahnärzteblatt* 2002;**42**(3):3.
- Rodemann 1999**  
Rodemann HP, Hehr T, Bamberg M. Relevance of the radioprotective effect of sodium selenite [Relevanz der radioprotektiven Wirkung von Natriumselenit]. *Medizinische Klinik* 1999;**94**(Suppl III):39–41.
- Sagowski 2004**  
Sagowski C, Wenzel S, Tesche S, Jenicke L, Kehrl W, Roeser K, Metternich FU. Sodium selenite reduces acute radiogenic damage of the rat parotid glands during fractionated irradiation [Reduktion akuter radiogener schädigungen durch natriumselenit. Wirksamkeit unter fraktionierter bestrahlung der glandula parotis der ratte]. *HNO* 2004;**52**(12):1067–75.
- Schleicher 1999**  
Schleicher UM, Lopez Cotarelo C, Andreopoulos D, Handt S, Ammon J. Radioprotection of human endothelial cells by sodium selenite [Radioprotektion humaner endothelzellen durch natriumselenit]. *Medizinische Klinik* 1999;**94**(Suppl. III):35–8.
- See 2006**  
See KA, Lavercombe PS, Dillon J, Ginsberg R. Accidental death from acute selenium poisoning. *The Medical Journal of Australia* 2006;**185**(7):388–9.
- Sehouli 2000**  
Sehouli J, David M, Kaufmann B, Lichtenegger W. Unconventional methods in cancer medicine - postoperative use by patients with gynecologic malignomas [Unkonventionelle methoden in der krebsmedizin – postoperative nutzung durch patientinnen mit gynäkologischen malignomen]. *Geburtshilfe Frauenheilkunde* 2000;**60**:146–54.
- Sill-Steffens 2003**  
Sill-Steffens R. Selenium in oncology [Selen in der onkologie]. *Deutsche Zeitschrift für Onkologie* 2003;**35**: 112–22.
- Trotti 2003**  
Trotti A, Bellm LA, Epstein JB, Frame D, Fuchs HJ, Gwede CK, et al. Mucositis incidence, severity and associated outcomes in patients with head and neck cancer receiving radiotherapy with or without chemotherapy: a systematic literature review. *Radiotherapy & Oncology* 2003;**66**(3): 253–62.
- Verhagen 1998**  
Verhagen AP, De Vet HC, De Bie RA, Kessels AGH, Boers M, Boeter LM, et al. The Delphi List: A criteria list for quality assessment of randomized clinical trials for conducting systematic reviews developed by delphi consensus. *Journal of Clinical Epidemiology* 1998;**51**(12): 1235–41.

**Vinceti 2001**

Vinceti M, Wei ET, Malagoli C, Bergomi M, Vivoli G. Adverse health effects of selenium in humans. *Reviews in Environmental Health* 2001;**16**(4):233–51.

**Weijl 1997**

Weijl NI, Cleton FJ, Osanto S. Free radicals and antioxidants in chemotherapy-induced toxicity. *Cancer Treatment Reviews* 1997;**23**(4):209–40.

**Yoshida 2000**

Yoshida M, Iizuka K, Terada A, et al. Prevention of nephrotoxicity of cisplatin by repeated oral administration of ebselen in rats. *The Tohoku Journal of Experimental Medicine* 2000;**191**:209–20.

**Zimmermann 1997**

Zimmermann T, Albrecht S, Kuhne H, Vogelsang U, Grutzmann R, Kopprasch S. Selenium administration in patients with sepsis syndrome. A prospective randomized study [Selensubstitution bei Sepsispatienten]. *Medizinische Klinik* 1997;**92** Suppl 3:3–4.

**References to other published versions of this review****Dennert 2006**

Dennert G, Horneber M. Selenium for alleviating the side effects of chemotherapy, radiotherapy and surgery in cancer patients. *Cochrane Database of Systematic Reviews* 2006, Issue 3. [DOI: 10.1002/14651858.CD005037.pub2

\* Indicates the major publication for the study

## CHARACTERISTICS OF STUDIES

### Characteristics of included studies [ordered by study ID]

#### Kasseroller 1998

Methods	Randomised placebo-controlled, double-blind trial with two parallel arms Recruitment period: July 1994 - September 1995 Observation period: 15 weeks Ethical approval: yes	
Participants	Number of patients: eligible/accrued: unclear - included: 60 - reported: 57 Drop Outs: "During the treatment, some of the patients were excluded from the study since they did not meet the criteria for study inclusion, namely, too short a period of stay." (Kasseroller 1998, p. 2228) Inclusion criteria: female cancer patients with one-sided secondary lymphoedema (arm) after mastectomy with axillar dissection and recurrent erysipela infections (> = 3 in the year before admission) who were admitted to the hospital for lymphoedema treatment Exclusion criteria: current treatment with antibiotics, NSAIDs or antioxidant medication Demographics: women, average age = 60.5 years Recruitment and setting: patients admitted to Wittlinger' s therapy center, Walchsee, Austria (private rehab clinic) for lymphoedema treatment Informed consent: yes	
Interventions	Intervention: week 1: sodium selenite solution 1000 µg/d p.o.; week 2-3: 300 µg/d p.o.; week 4-15: 200 µg/d p.o. (body weight > 70 kg), 100 µg/d p.o. (body weight < 70 kg) Control: identically looking placebo (sodium chloride p.o.) Concomitant treatment: combined physical congestion-relief program (manual lymph drainage twice daily, compressive bandages, high-voltage therapy, therapeutic exercise, meticulous skin care)	
Outcomes	Primary outcome measure: incidence of recurrent erysipelas in a 15-week follow-up period Outcome assessment: diagnosis of erysipela infection based on clinical examination, blood sedimentation rate and CRP titer	
Notes	Post-marketing surveillance study in Austria Participants: no data on stage of lymphoedema, comorbidity given Number of participants: inconsistent data: results reported for n = 57 participants in text, but for n = 60 participants in table Intervention: application of 1000 mg sodium selenite reported (misprint instead of 1000 µg) Outcomes: criteria for erysipela diagnosis unclear, diagnosis might differ between in-hospital treatment and follow-up after dismissal Investigator provided additional information on study methods	
<b><i>Risk of bias</i></b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>

**Kasseroller 1998** (Continued)

Allocation concealment (selection bias)	Unclear risk	B - Unclear
---	--------------	-------------

**Mücke 2007**

Methods	Ongoing randomised placebo-controlled, open label trial with 2 parallel arms Recruitment period: December 2000 - ongoing Observation period: December 2000 - ongoing most recent publication reports on December 2000 to May 2007 Ethical approval: yes
Participants	Number of patients eligible: unclear - blood tested for selenium deficiency: unclear - selenium deficiency in blood: unclear - included: unclear reported: 82 Drop Outs: unclear Inclusion criteria: female cancer patients with cervical or endometrial cancer after surgery and with pre-existing selenium deficiency in whole blood/serum receiving adjuvant percutaneous radiotherapy Exclusion criteria: distant metastases; patients receiving combined radio-/ chemotherapy; patients with prior pelvic/ abdominal radiation; current diarrhea; use of antibiotics, antacids, astringents, adsorbents, intestinal disinfectants or 5-HT3-antagonists at the beginning of the study; use of selenium supplements; use of cytoprotectants Demographics: women, age 31 to 80 years (median: 66 years). Recruitment and setting: multicenter study, Germany Informed consent: yes
Interventions	Intervention: sodium selenite solution 500 µg/d for two days before start of radiotherapy, 500 µg/d on days of radiation, 300 µg/d on days without radiation (weekend) Control: no supplementation all patients: percutaneous adjuvant radiotherapy with total doses between 45 to 50 Gy and single doses between 1.8 to 2.0 Gy supportive medication: not reported
Outcomes	Primary outcome measure: compensation of selenium deficiency as measured in whole blood other: incidence and severity of diarrhoea Outcome assessment: weekly examination by physician; blood tests; participants' diary of well-being and physical problems
Notes	Post-marketing surveillance study in Germany Intervention: concomitant medication with influence on diarrhea (opioids etc.) not reported Investigators provided a study plan and additional information on study methods Actual intervention deviated from study protocol: sodium selenite prior to radiotherapy was omitted in a number of patients (see: recent publications, confirmed by investigators)

*Risk of bias*

Mücke 2007 (Continued)

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

Zimmermann 2005

Methods	Randomised placebo-controlled, double-blind trial with 2 parallel arms Recruitment period: unclear Observation period: 2 weeks (for postoperative face swelling) Ethical approval: yes
Participants	Number of patients: eligible: unclear - included: unclear - reported: 20 Drop outs: unclear Inclusion criteria: female and male patients, age >18, with carcinoma of the tongue or floor of the mouth (clinical stage T3 or T4) Exclusion criteria: clinical tumour stage T1 or T2; second malignancy; cardiac oedema; hypoproteinic oedema; acute infections during the past 6 months; regular intake of antioxidants; chronic renal failure; intellectual disability; pregnant women; prison inmates; participants of other clinical studies Demographics: 18 men - 2 women; age 30 - 74 (mean: 57) years; all patients had histologically confirmed squamous cell cancer (2 pT2, 7 pT3, 8 pT4) Recruitment and setting: monocenter study, Dresden, Germany Informed consent: yes
Interventions	Intervention: 3000 µg sodium selenite solution i.v. on day of surgery; 1000 µg sodium selenite solution i.v. or p.o. on days 1 to 21 after surgery Control: placebo (0.9% sodium chloride solution) all patients: surgical tumour resection in curative intention with bilateral neck dissection
Outcomes	Primary outcome measures: three defined distances and circumferences in the face; selenium levels in whole blood and several other laboratory parameters Assessment of outcome of interest: manual measurement by one investigator at four time points (prior to, immediately after, one week after and two weeks after surgical intervention)
Notes	Intervention: concomitant surgical and non-surgical interventions with influence on lymphoedema not reported Investigators provided additional information on their interpretation of study results All patients who refused the continuation of study medication due to perceived adverse effects would have been excluded from the final study report according to Leonhardt 2002; according to personal communication no drop-outs occurred

*Risk of bias*

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Unclear risk	B -Unclear

CRP: C-reactive protein  
d: day  
Gy: Gray  
µg: microgram  
NSAID: non-steroidal antiinflammatory drugs  
p.o.: per os

### Characteristics of excluded studies *[ordered by study ID]*

Study	Reason for exclusion
<a href="#">Asfour 2006</a>	RCT, outcome:apoptosis in polymorphonuclear leucocytes; also infections in selenium and control group reported, authors contacted. Study excluded because assessment is not possible on the basis of available information
<a href="#">Badger 2004</a>	Cochrane Review: includes patients with malignant and non-malignant diseases
<a href="#">Büntzel 1999</a>	Case series
<a href="#">Elango 2006</a>	Clinical trial; outcome:plasma selenium level and activity of selenium-dependent enzymes
<a href="#">Fakih 2006</a>	Phase I clinical trial to determine the maximum tolerated dose of irinotecan in combination with a fixed dose of selenomethionine
<a href="#">Federico 2001</a>	RCT, zinc/selenium combination supplement
<a href="#">Fraunholz 2001</a>	Non-randomized clinical trial
<a href="#">Hoenjet 2005</a>	RCT with multi component supplement; only non-clinical outcomes
<a href="#">Hu 1997</a>	RCT, outcome: treatment toxicities of chemotherapy; data not presented according to toxicity scale; author contacted, but no reply obtained
<a href="#">Kasseroller 1995</a>	RCT, participants with malignant and non-malignant diseases
<a href="#">Kiremidjian 2000</a>	RCT, outcome: reaction of peripheral lymphocytes to mitogen stimulation after selenium supplementation
<a href="#">Kreienberg 2003</a>	Study title mentioned in University of Heidelberg´ s trials´ register of the year 2003; hospital contacted, but no answer obtained
<a href="#">Lasch 1999</a>	Clinical controlled trial: study not described as randomized, allocation to study and control group was performed according to date of birth
<a href="#">Schumacher 2003</a>	Planned RCT, so far not recruiting patients
<a href="#">Sieja 2000</a>	Multi component drug, study not described as randomised
<a href="#">Sundström 1984</a>	No RCT, outcome: change in selenium-related biomarkers

(Continued)

<a href="#">Sundström 1989</a>	No RCT, outcome: change in selenium-related biomarkers
<a href="#">Wagler 2000</a>	RCT, outcome: change in selenium-related biomarkers - author contacted, but no reply obtained
<a href="#">Xu 1990</a>	RCT, outcome: change of chemiluminescence measurement after selenium supplementation
<a href="#">Xu 1999</a>	RCT, outcome: immunomodulation following selenium supplementation
<a href="#">Yu 1996</a>	Outcome: selenium-related biomarkers; author contacted, but no answer obtained

### Characteristics of ongoing studies [ordered by study ID]

#### [Büntzel 2004](#)

Trial name or title	Einsatz von selenase als Zusatztherapie zur Radiotherapie bei Tumoren im Hals-Nasen-Ohrenbereich. [Use of selenase as complementary therapy for radiotherapy in ENT tumours]
Methods	
Participants	200 female and male patients with selenium deficiency in whole blood receiving radiotherapy for ENT carcinomas
Interventions	Supplementation of sodium selenite solution p.o. Intervention: 500 µg/d for two days before start of radiotherapy, 500 µg/d on days of radiation, 300 µg/d on days without radiation (weekend) Control: no supplementation all patients: percutaneous adjuvant radiotherapy with total doses between 45 to 50 Gy and single doses between 1.8 and 2.0 Gy
Outcomes	Primary outcome measure: normalisation of pre-existing selenium deficiency in whole blood other: radiotherapy toxicities
Starting date	01.09.2000
Contact information	Dr Jens Büntzel, Südharzkrankenhaus Nordhausen, Klinik für HNO-Erkrankungen, Dr.-Robert-Koch-Str. 29, 99734 Nordhausen, Germany
Notes	Information taken from study protocol version 1.4/01.03.2004 Study is recruiting according to authors

## DATA AND ANALYSES

This review has no analyses.

## ADDITIONAL TABLES

Table 1. Quality assessment of included studies

Study	Allocation	Concealment	Study groups	Eligibility	Blinding	Attrition	Delphi list	Jadad score	Comment
<a href="#">Kasseroller 1998</a>	random allocation; method of sequence generation not reported	unclear	no information on comparability of study groups regarding risk factors at baseline	eligibility criteria specified; number of eligible patients not stated	patient: yes; care provider: yes; outcome assessor: yes - unblinding of treatment status after end of intervention	3 dropouts	1-0-0-1-1-1-1-0-0	1-1-1	Selection bias possible: allocation concealment unclear, no information on distribution of risk factors at baseline. Observer/detection bias possible: no control procedure reported whether blinding was successful, details on outcome assessment not reported. Performance bias possible: concomitant treatment not sufficiently described
<a href="#">Mücke 2007</a>	random allocation; method of	serially numbered, sealed,	no information on compa-	eli- gibility cri- teria speci-	patient: no; care provider:	dropouts / with-	--	--	Preliminary

**Table 1. Quality assessment of included studies** (Continued)

	sequence generation not reported	opaque envelopes	reliability of study groups regarding risk factors at baseline	defined; number of eligible patients not stated	no; outcome assessor: no	drawals not reported in most recent publications			results of an ongoing study (Delphi List and Jadad Score will be given when final report is available) Selection bias unlikely, but possible: allocation concealment adequate, but no information on distribution of risk factors for outcome. Observer/detection bias possible: no blinding. Performance bias possible: concomitant treatment unclear
Zimmermann 2005	random allocation; method of sequence generation not reported	unclear	no information on comparability of study groups regarding risk factors at baseline	eligibility criteria specified; number of eligible patients not stated	patient: unclear; care provider: unclear; outcome assessor: unclear	no dropouts or withdrawals	1-0-0-1-0-0-0-0-0	1-1-1	Selection bias possible: allocation concealment unclear, information on distribution of

**Table 1. Quality assessment of included studies** (Continued)

										<p>risk factors at baseline</p> <p>Observer/detection bias possible: study was reported as double-blinded - informed consent form, however, contained information on intervention status, reliability and validity of outcome assessment unclear</p> <p>Performance bias possible: concomitant treatment not sufficiently described</p> <p>Randomisation and blinding unclear: informed consent form for participants contains the sentence</p>
--	--	--	--	--	--	--	--	--	--	---

**Table 1. Quality assessment of included studies** (Continued)

										“you have been allocated to the group receiving selenium supplements”, while study is reported as double-blinded
--	--	--	--	--	--	--	--	--	--	--

## APPENDICES

### Appendix I. MEDLINE search strategy

1. selen\* OR seleno\* OR selenium[mesh] OR selenium compounds[mesh] OR organoseleno compounds[mesh]
2. epidemiologic study characteristics[mesh] OR random\* OR placebo\* OR clinical trial\* OR double blind method OR single blind method
3. neoplasms[mesh] OR stem cell transplantation[mesh] OR therapeutics[mesh] OR antineoplastic agents[mesh] OR quality of life[mesh]
4. cancer\* OR malignan\* OR carcino\* OR neoplasm\* OR tumor\* OR tumour\*
5. radioth\* OR radiat\* OR irradiat\* OR radiochemo\* OR radio-chemo\* OR chemotherap\*
6. melanoma\* OR sarcoma\* OR adenoma\* OR adenosarcoma\* OR adenocarcinoma\* OR carcinosarcoma\* OR chondrosarcoma\* OR fibrosarcoma\* OR dermatofibrosarcoma\* OR neurofibrosarcoma\* OR hemangiosarcoma\* OR leiomyosarcoma\* OR liposarcoma\* OR myosarcoma\* OR rhabdomyosarcoma\* OR myxosarcoma\* OR osteosarcoma\* OR lymphoma\* OR stem cell transplant\* OR stem-cell transplant\* OR bone marrow transplant\*
7. lymphedem\* OR lymphoedem\* OR cardiotox\* OR nausea\* OR vomit\* OR hair loss\* OR leukopenia\* OR neutropenia\* OR thrombopenia\* OR anemia\*
8. #3 OR #4 OR #5 OR #6 OR #7
9. #1 AND #2 AND #8
10. animals[mh] NOT humans[mh]
11. #9 NOT #10

## Appendix 2. Additional search strategies

Database	Time	Search Strategy
EMBASE	1980 - 2007 week 28	1 selenium/ or selen\$.mp. 2 exp Selenium Derivative/ 3 methylseleninic acid/ or methylselenium.mp. 4 exp Organoselenium Derivative/ 5 exp Clinical Study/ 6 exp NEOPLASM/ 7 Selenium 75/ 8 1 or 2 or 3 or 4 9 8 and 5 and 6 10 9 not 7 11 limit 10 to human
SIGLE [ <a href="http://www.fiz-informationsdienste.de">www.fiz-informationsdienste.de</a> ] [the database was discontinued]	October 2004	?selen? with restriction to subject group "06 Biological and Medical Sciences (without Botany, Zoology, Biophysics)"
Cancerlit	July 2007	1 random* OR placebo* OR clinical trial* OR randomized controlled trial* OR controlled clinical trial* OR double blind OR single blind 2 selen* OR organoselen* 3 1 AND 2
Clinical Contents in Medicine CCMed	October 2004	selen* OR organoselen* OR natriumselen*
NCI Clinical Trials [ <a href="http://www.cancer.gov">www.cancer.gov</a> ]	October 2004	medication=selenium
ISRCTN and mRCT [ <a href="http://www.controlled-trials.com">www.controlled-trials.com</a> ]	October 2004	selenium AND cancer
German Register of Cancer Studies [ <a href="http://www.studien.de">www.studien.de</a> ]	July 2007	selen
Cochrane Pain, Palliative & Supportive Care Trials Register	July 2007	selen*
The Cochrane Database of Systematic Reviews; The Database of Abstracts of Reviews of Effects (DARE); The Cochrane Central Register of Controlled Trials (CENTRAL)	Issue 2, 2007	selen*

## WHAT'S NEW

Date	Event	Description
16 March 2017	Review declared as stable	See <a href="#">Published notes</a> .

## HISTORY

Date	Event	Description
11 May 2011	Amended	Contact details updated.
8 February 2011	Amended	Contact details updated.
24 September 2010	Amended	Contact details updated.
17 February 2009	New search has been performed	For this update review one new included study ( <a href="#">Zimmermann 2005</a> ) was included adding a further 20 participants of head and neck cancer patients to the total participants included within the review. The <a href="#">Mücke 2007</a> latest data included a further 19 participants. The previous review identified an ongoing study which has been completed since the original publication of this review and is now excluded ( <a href="#">Schumacher 2003</a> ). The list of excluded studies, background and discussion chapter was also revised. The search for this update was run in July 2007. The new included and excluded studies did not change the original conclusions of this review
22 April 2008	Amended	Converted to new review format.

## CONTRIBUTIONS OF AUTHORS

GD, MH: protocol development, study assessment, data extraction, data analysis, discussion.

GD: literature search, data management, review coordination, contact with first authors, writing, review update.

## DECLARATIONS OF INTEREST

None known

## SOURCES OF SUPPORT

### Internal sources

- No sources of support supplied

### External sources

- Dr Ernst and Anita Bauer Stiftung, Germany.
- EU Project: Concerted action for complementary and alternative medicine in the cancer field (EU CAM-Cancer) (contract no.: QLG4-CT-2002-00786), Other.

## NOTES

We performed a full search in September 2016 and did not identify any potentially relevant studies likely to change the conclusions. Therefore, this review has now been stabilised following discussion with the authors and editors. If appropriate, we will update the review if new evidence likely to change the conclusions is published.

During planning for the potential update, the author team was amended, and we acknowledge their contribution: Peter Renner<sup>1</sup>, Alexander Kalisch<sup>2</sup>.

<sup>1</sup>Department of Internal Medicine, Division of Cardiology, Paracelsus Medical University, Klinikum Nuremberg, Nuremberg, Germany

<sup>2</sup>Department of Internal Medicine, Division of Oncology and Hematology, Paracelsus Medical University, Klinikum Nuremberg, Nuremberg, Germany

## INDEX TERMS

### Medical Subject Headings (MeSH)

\*Neoplasms [drug therapy; radiotherapy; surgery]; Antineoplastic Agents [adverse effects]; Antioxidants [\*therapeutic use]; Diarrhea [\*prevention & control]; Lymphedema [\*prevention & control]; Postoperative Complications [prevention & control]; Radiotherapy [adverse effects]; Randomized Controlled Trials as Topic; Sodium Selenite [\*therapeutic use]

### MeSH check words

Humans